

Adeleks



Adeleks tablet 4-8 mg, 20 tablets Adeleks IM 4 mg/2 mL ampoule

COMPOSITION: Each tablet contains 4-8 mg thiocolchicoside. Each ampoule contains 4 mg thiocolchicoside, 16.8 mg sodium chloride and q.s. 2 mL water for injection.

PHARMACOLOGICAL PROPERTIES: Thiocolchicoside, the active ingredient of Adeleks, is a thio-methyl derivative of colchicoside which is a colchicine glucoside. Thiocolchicoside has a myorelaxant effect on muscles and especially on striated muscles.

INDICATIONS: Adeleks is indicated in symptomatic treatment of painful muscle cramps.

CONTRAINDICATIONS: Adeleks should not be used in individuals hypersensitive to thiocolchicoside or colchicine, in hypotonic muscle conditions and in loose paralysis. Intramuscular administration is contraindicated in patients receiving anti-coagulant therapy and in patients with bleeding.

WARNINGS AND PRECAUTIONS: Thiocolchicoside should not be used in patients with severe renal failure. Effect on driving and operating machinery: Care should be exercised when driving and operating machinery. Use in pregnancy: **Pregnancy category is X.**

SIDE EFFECTS / ADVERSE EFFECTS: Gastrointestinal system side effects of gastralgia and diarrhea can be observed during thiocolchicoside administration. Although rarely, decrease in blood pressure, temporary loss of consciousness or excitation can be seen. During thiocolchicoside therapy, rash can develop in hypersensitive patients. Consult your doctor in case of an unexpected effect.

DRUG INTERACTIONS: There is no data on drug interactions of thiocolchicoside. However, when administered concomitantly with similar drugs, the patient should be closely monitored.

DOSAGE AND ADMINISTRATION: ADELEKS tablets should be taken on a full stomach. Unless prescribed otherwise by the physician, ADELEKS 4 mg tablet should be given 2 tablets 2 times a day and ADELEKS 8 mg tablet should be given 1 tablet (max. dosage 16 mg a day) 2 times a day bid for severe muscle cramps. Use in children under the age of 15 years is not recommended. Unless recommended otherwise by the physician, ADELEKS IM ampoule should be administered bid via intramuscular route in 2 mg doses. It is recommended that IM therapy should be continued for 3-5 days. If the treatment is to be continued further, therapy should be switched to oral administration and continued by 2 tablets bid for 5-7 days. An effect period of 30-40 minutes should lapse for physical therapy sessions. Use in children under the age of 15 years is not recommended.

STORAGE CONDITIONS: It should be kept out of reach of children, in its package and in room temperature <25°C.

AVAILABLE FORMS:

ADELEKS 4-8 mg tablet in 20 tablets blister packs.

ADELEKS 4 mg / 2 mL IM ampoule, in 6-ampoules packages.

NAME AND ADDRESS OF THE AUTHORISATION HOLDER: Mustafa Nevzat İlaç Sanayii A.Ş.
Prof. Dr. Bülent Tarcan Sok., Pak İş Merkezi No: 5/1 34349 Gayrettepe/İstanbul.

NAME AND ADDRESS OF THE MANUFACTURING COMPANY: Mustafa Nevzat İlaç Sanayii A.Ş. Çobançeşme Mah. Sanayi Cad. No:13 Yenibosna-İstanbul.