

Oksamen



Oksamen – L freeze-dried for injection 20 mg, 1 vial + 1 diluent ampoule
Oksamen film-coated tablet 20 mg, 10 film-coated tablet

COMPOSITION: Each film-coated tablet contains 20 mg tenoxicam.
Each freeze-dried injection contains 20 mg tenoxicam.

PHARMACOLOGICAL PROPERTIES: Has antiinflammatory, analgesic, antipyretic and anti-rheumatic activity.

INDICATIONS:

Rheumatoid arthritis
Osteoarthritis
Ankylosing spondylitis
Arthrosis

Non-articular conditions such as tendinitis, bursitis, peri-arthritis of the shoulder or hips, sprains and strains

Painful, inflammatory and degenerative diseases of the musculoskeletal system such as epicondylitis, sciatica, lumbago, acute gout and traumatic cases

CONTRAINDICATIONS: Is contraindicated in patients who are hypersensitive to tenoxicam and also in patients who have a history of gastric or duodenal ulcer, gastrointestinal bleeding and chronic gastritis. Should not be administered in patients with severe renal failure or hepatic insufficiency.

WARNINGS AND PRECAUTIONS: Warnings and precautions required for all non-steroidal antiinflammatory drugs are also valid for OKSAMEN tablet and OKSAMEN-L for injection.

SIDE EFFECTS / ADVERSE EFFECTS: In prolonged clinical trials with tenoxicam (1 year) the drug was well tolerated when given in the recommended dosage. Side effects that may be seen with OKSAMEN-L for injection and OKSAMEN tablet are listed below: Gastrointestinal tract, Central nervous system, Skin and soft tissue, Urinary tract, Liver and biliary tract.

DRUG INTERACTIONS: Tenoxicam, at therapeutic doses, has no pharmacokinetic interaction with antacids, cimetidine, oral antidiabetics and oral anticoagulant agents; however, it can increase the effects of anticoagulant agents such as coumarine.

DOSAGE AND ADMINISTRATION: The recommended dose for OKSAMEN tablet is 20 mg once daily. In prolonged therapies, maintenance dose is 10 mg daily. The recommended dose for OKSAMEN-L for injection is 20 mg once daily. In the treatment of acute gout arthritis, OKSAMEN-L for injection or OKSAMEN tablet is administered 40 mg daily for the first 2 days and 20 mg daily for the following 5 days.

STORAGE CONDITIONS: Should be kept out of reach of children, at room temperature (< 25 °C) and in its package.

AVAILABLE FORMS: OKSAMEN tablet 20 mg, in 10-tablet blister packs

OKSAMEN-L freeze-dried for injection 20 mg, 1 vial + 1 diluent ampoule (2 mL)

NAME AND ADDRESS OF THE AUTHORISATION HOLDER: Mustafa Nevzat İlaç Sanayii A.Ş.
Prof. Dr. Bülent Tarcan Sok., Pak İş Merkezi No: 5/1 34349 Gayrettepe/İstanbul.

NAME AND ADDRESS OF THE MANUFACTURING COMPANY: Mustafa Nevzat İlaç Sanayii A.Ş. Çobançeşme Mah. Sanayi Cad. No:13 Yenibosna-İstanbul.